

The Effect of Accumulation of Excess Energy of a Body in Gravitational Compression

S. S. Gershtein, A. A. Logunov, M. A. Mestvirishvili
Institute for High Energy Physics, Protvino, Russia

Abstract

It is shown that a gravitational compression of a spherical body results in an infinite growth of the energy of a body as its radius comes close to GM/c^2 . This gives rise to a negative defect of mass and, due to an instability, to an expansion or to an explosion. A rigorous proof of the above statement can be obtained within the General Relativity in the harmonic coordinates.

The Hilbert-Einstein equation in arbitrary coordinates has the form [1]

$$(-g)(T^{ik} + t_g^{ik}) = \frac{\partial h^{ikl}}{\partial x^l}, \quad (1)$$

$$\text{where } h^{ikl} = \frac{\partial}{\partial x^m} \lambda^{iklm}, \quad \lambda^{iklm} = \frac{1}{16\pi} (-g)(g^{ik} g^{lm} - g^{il} g^{km}). \quad (2)$$

We use the system of units in which $G = c = 1$. From formula (2) it follows that

$$h^{ikl} = -h^{ilk}. \quad (3)$$

Now, Eqs. (1) and (3) imply the conservation law

$$\frac{\partial}{\partial x^k} (-g)(T^{ik} + t_g^{ik}) = 0, \quad (4)$$

which is valid in an arbitrary reference frame. Here t_g^{ik} is the Landau-Lifshitz pseudotensor [1]:

$$\begin{aligned} (-g)t_g^{ik} = & \frac{1}{16\pi} \{ \partial_l \tilde{g}^{ik} \partial_m \tilde{g}^{lm} - \partial_l \tilde{g}^{il} \partial_m \tilde{g}^{km} + \frac{1}{2} \tilde{g}^{ik} \tilde{g}_{lm} \partial_p \tilde{g}^{ln} \partial_n \tilde{g}^{pm} - \\ & - (\tilde{g}^{il} \tilde{g}_{mn} \partial_p \tilde{g}^{kn} \partial_l \tilde{g}^{mp} + \tilde{g}^{kl} \tilde{g}_{mn} \partial_p \tilde{g}^{in} \partial_l \tilde{g}^{mp}) + \\ & + \tilde{g}_{lm} \tilde{g}^{np} \partial_n \tilde{g}^{il} \partial_p \tilde{g}^{km} + \frac{1}{8} (2\tilde{g}^{il} \tilde{g}^{km} - \tilde{g}^{ik} \tilde{g}^{lm}) (2\tilde{g}_{np} \tilde{g}_{qr} - \tilde{g}_{pq} \tilde{g}_{nr}) \partial_l \tilde{g}^{nr} \partial_m \tilde{g}^{pq} \}, \end{aligned} \quad (5)$$

and $\tilde{g}^{mn} = \sqrt{-g} g^{mn}$; $m, n = 0, 1, 2, 3$. It should be noted that, in the Relativistic Theory of Gravitation (RTG), the conservation law (4) is valid only in an inertial reference frame in the Galilean coordinates. In what follows, we use the pseudotensor for calculations in General Relativity (GR) only in the harmonic coordinates. In view of formula (1), the Landau-Lifshitz pseudotensor outside of a body coincides with the Fock pseudotensor and formula (4) implies conservation of the following integrals of motion [1]:

$$P^i = \int (-g)(T^{ik} + t_g^{ik}) dS_k. \quad (6)$$

Performing an integration over the hypersurface $x^0 = \text{const}$, we arrive at the integral over three-dimensional space,

$$P^i = \int (-g)(T^{i0} + t_g^{i0})dV. \quad (7)$$

Our considerations rely heavily on the Hilbert causality principle, however, it receives only a little attention in the literature. For this reason, we give an extended quotation from the Hilbert paper [2]:

“So far we considered all coordinate systems x_s obtained from some system by an arbitrary coordinate transformations as equivalent. This arbitrariness must be limited if we consider that two points located on the same time line remain in casual relationship to each other and, therefore, they cannot go over into two equal-time points under a proper coordinate transformation.

Assuming that x_0 is the **proper-time coordinate**, we formulate the following definition. A system of space-time coordinates is named a **proper** coordinate system if, in addition to the condition $g < 0$, the inequalities

$$g_{11} < 0, \quad \begin{vmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{vmatrix} > 0, \quad \begin{vmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \\ g_{31} & g_{32} & g_{33} \end{vmatrix} < 0, \quad g_{00} > 0. \quad (\alpha)$$

are valid. *A transformation which maps a proper system of space-time coordinates into a proper system of space-time coordinates is named a **proper** coordinate transformation”. Some later Hilbert notes: “Since under a proper coordinate transformation the time line goes over into time line, two world points located at the same time line never go over into two world points with the same value of time coordinate; that is, they cannot be associated with simultaneous events”... “In the gravitational field, a single mass point moves along geodesic, which represents a time line. Light moves along a null geodesic.*

*A world line, which is a trajectory of a test particle, must be a time line. For this reason, we can always go over into rest frame of this particle using **proper** coordinate transformations”.*

The Hilbert conditions (α) on the metric coefficients are indispensable.

Hilbert emphasized this: *“We see that the fundamental concept of cause and effect underlying the causality principle does not give rise to inconsistencies when we add the inequalities (α) to our main equations, that is, if we restrict our consideration to the **proper** space-time coordinates”.*

Hilbert causality conditions keep the metric signature $(+ - - -)$ invariant under proper coordinate transformations. From the inequalities (α) it is seen that the Hilbert causality conditions put no limit on the metric coefficients $g_{0\alpha}$.

The Hilbert causality principle is of fundamental importance. This means that only those solutions of the Hilbert-Einstein equations make physical sense in General Relativity, which satisfy the Hilbert causality conditions.

It was shown [3] that the GR equations with regard of the Hilbert causality principle rule out a possibility of the gravitational collapse. However, behavior of the radius $r_0(t)$ of a body in gravitational compression has not received sufficient attention. In this work, we study its behavior by considering the integral of motion in harmonic coordinates, which are privileged coordinates as it was shown by academician V.A. Fock.

In the GR, in the case of non-static spherically symmetric body, an interval outside of the body is static. In the harmonic coordinates it has the form [4]

$$ds^2 = \frac{r-M}{r+M}dt^2 - \frac{r+M}{r-M}dr^2 - (r+M)^2(d\theta^2 + \sin^2\theta d\phi^2). \quad (8)$$

Changing from the spherical to the Cartesian coordinates, we arrive at $(\alpha, \beta = 1, 2, 3)$

$$g_{00} = \frac{r-M}{r+M}, \quad g_{0\alpha} = 0, \quad g_{\alpha\beta} = -(1 + \frac{M}{r})^2 \delta_{\alpha\beta} - \frac{r+M}{r-M} \frac{M^2}{r^4} x_\alpha x_\beta. \quad (9)$$

From these expressions we obtain [4]

$$\sqrt{-g}g^{\alpha\beta} = \tilde{g}^{\alpha\beta} = -\delta_{\alpha\beta} + M^2 \frac{x_\alpha x_\beta}{r^4}, \quad \tilde{g}^{00} = \frac{(r+M)^3}{r^2(r-M)}, \quad \tilde{g}^{0\alpha} = 0. \quad (10)$$

From the above formulas it follows that the coordinates x_α are harmonic [4],

$$\frac{\partial \tilde{g}^{\alpha\beta}}{\partial x^\beta} = 0. \quad (11)$$

Following the ideas of paper [5], we rearrange the integral of motion (7) to the form

$$P^0 = \int_{V_0(t)} (-g)(T^{00} + t_g^{00})dV + \int_{V_1(t)} (-g)t_g^{00}dV, \quad (12)$$

where $V_0(t)$ is the interior of the body and $V_1(t)$ is the (infinite) domain outside the body. An integration over the domain $V_1(t)$ with the use of harmonic coordinates x_α gives

$$\int_{V_1(t)} (-g)t_g^{00}dV = -M^2 \left(\frac{4}{r_0(t) - M} - \frac{1}{2r_0(t)} + \frac{M}{r_0^2(t)} + \frac{M^2}{2r_0^3(t)} \right). \quad (13)$$

The integral of motion (7) is readily calculated by reducing to an integral over a surface [1]

$$P^0 = M. \quad (14)$$

Substituting expressions (13) and (14) in formula (12), we obtain

$$\int_{V_0(t)} (-g)(T^{00} + t_g^{00})dV = M + M^2 \left[\frac{4}{r_0(t) - M} - \frac{1}{2r_0(t)} + \frac{M}{r_0^2(t)} + \frac{M^2}{2r_0^3(t)} \right]. \quad (15)$$

This formula was derived earlier in the Relativistic Theory of Gravitation (RTG) due to the existence of the energy-momentum *tensor* of the gravitational field [5]. The physical solution is obtained by "sewing" of the non-static spherically symmetric solution inside of a body with the solution (8) in the exterior domain. Since it satisfies the Hilbert causality conditions, the volume of a body in gravitational compression is always finite and, therefore, the energy of a body is also finite [6].

Finiteness of the energy of a body in gravitational compression, as it follows from formula (15), rules out a possibility of gravitational collapse. Thus we conclude that the quantity GM/c^2 is of fundamental importance — it gives the radius that cannot be approached by a body of mass M :

$$r_0(t) > M. \quad (16)$$

Therefore, a massive body cannot be pointlike. A calculation of the value of $t_g^{ik} g_{ik} = t_g$ in the GR in harmonic coordinates gives [5]

$$32\pi t_g = -\frac{r_g^2(24r^2 - 16rr_g + 5r_g^2)}{4r^2(r - M)(r + M)^3}; \quad r_g = 2M. \quad (17)$$

The value of t_g must be limited in order to sew solutions in the interior and exterior domains, i.e. it is necessary

$$r > M, \quad (18)$$

in agreement with the condition (16). In RTG, the quantity t_g is an invariant of the gravitational field [5].

From formula (15) it follows that, when thermal and thermonuclear sources of energy of a star are exhausted, a star of large mass ($M > 3M_\odot$) accumulates an excess of positive energy from gravitational compression, which results in a negative mass defect. This, in its turn, stops compression and gives rise to a subsequent radial expansion of a body. Due to an instability, such expansion can be of explosive type providing the validity of inequality (16). After multiple repetition of such process it may be that only stable objects remain.

Our conclusions agree with the empiric model proposed by Ambarzumian in 1960. In this model (see [7]), “superdense matter evolves into a diffuse state; positive-energy states can explode and transform from dense to diffuse state”. For more detail, special computations are needed. States with excess energy as well as other positive-energy solutions were considered in [7],[8].

Provided that the Landau–Lifshitz pseudotensor is calculated in the GR in harmonic coordinates, one obtains neither gravitational collapse nor black holes.

We are grateful to V.A. Petrov, A.P. Samokhin and N.E. Tyurin for useful discussions.

References

- [1] *L.D. Landau and E.M. Lifshitz*, Field Theory. Nauka, Moscow, 2001, p. 379, (in Russian).
- [2] *D. Hilbert*, Math. Ann., Bd. 92 (1924) S. 1-32.
- [3] *A.A. Logunov and M.A. Mestvirishvili*, Theor. Math. Phys., Vol. 170 (2012) pp. 413-419.
- [4] *V.A. Fock*, The Theory of Space, Time and Gravitation. Pergamon Press, 1964.
- [5] *A.A. Logunov and M.A. Mestvirishvili*, Theor. Math. Phys., Vol. 174 (2013) pp. 253-262.
- [6] *J.L. Synge*, Relativity: The General Theory. North-Holland Publ. Company, Amsterdam, 1960.
- [7] *Ya.B. Zel'dovich and I.D. Novikov*, Relativistic Astrophysics. Nauka, Moscow, 1967, p. 35, (in Russian).
- [8] *B.K. Harrison, K.S. Thorne, M. Wakano and J.A. Wheeler*, Gravitation theory and gravitational collapse. The University of Chicago Press, Chicago and London, 1965.